



## National Library Publ

Nucleotide	Protein	Genome	Structure	PopSet		
for <u>9924606</u>	3			Go Clear		
Limits	Preview/Ind	lex Histo	ory Clipbos	ırd		
Display Sun	nmary 🔻	Save Text	Order Details	Add to Clipboard		
1: Nagase T, Ishikawa K, Suyama M, Kikuno R, Hirosawa M, Miyajima N, Tanaka A, Kotani H, Nomura N, Ohara O.						
complete proteins i DNA Res.	e sequences of 10 in vitro. 1999 Feb 26;6(1):6	00 new cDNA 3-70.				
	Prediction complete proteins DNA Res.	for 99246063 Limits Preview/Inc  Display Summery  1: Nagase T, Ishikawa K, Suyama M, Miyajima N, Tanaka A, Ko O.  Prediction of the coding s complete sequences of 10 proteins in vitro. DNA Res. 1999 Feb 26;6(1):6	for 99246063  Limits Preview/Index History  Display Summary Save Text  1: Nagase T, Ishikawa K, Suyama M, Kikuno R, H, M, Miyajima N, Tanaka A, Kotani H, Nomura NO.  Prediction of the coding sequences of use complete sequences of 100 new cDNA	Nucleotide		

Related Resources

Write to the Help Desk
NCBI | NLM | NIH
Department of Health & Human Services
Freedom of Information Act | Disclaimer





## National Library PubMed of Medicine

PubMed	Nucleotide	Protein	Genome	Structure	PopSet
Search PubMed	▼ for			Go	Clear
·	Limits	Preview/In	dex History	Clipboard	I
	<b>*</b>				
•••	Display A	bstract: v	Save Text C	order Add to Cl	ipboard
Cataon Dublidad				***************************************	

Entrez PubMed

☐ 1: DNA Res 1999 Feb 26;6(1):63-70

Related Articles, Books, Protein, Nucleotide, LinkOut

**PubMed Services** 

Prediction of the coding sequences of unidentified human genes. XIII. The complete sequences of 100 new cDNA clones from brain which code for large proteins in vitro.

Nagase T, Ishikawa K, Suyama M, Kikuno R, Hirosawa M, Miyajima N, Tanaka A, Kotani H, Nomura N, Ohara O

Related Resources

Kazusa DNA Research Institute, Kisarazu, Chiba, Japan.

As a part of our cDNA project for deducing the coding sequence of unidentified human genes, we newly determined the sequences of 100 cDNA clones from a set of size-fractionated human brain cDNA libraries, and predicted the coding sequences of the corresponding genes, named KIAA0919 to KIAA1018. The sequencing of these clones revealed that the average sizes of the inserts and corresponding open reading frames were 4.9 kb and 2.6 kb (882 amino acid residues), respectively. A computer search of the sequences against the public databases indicated that predicted coding sequences of 87 genes contained sequences similar to known genes, 53% of which (46 genes) were categorized as proteins relating to cell signaling/communication, cell structure/motility and nucleic acid management. The chromosomal locations of the genes were determined by using human-rodent hybrid panels unless their mapping data were already available in the public databases. The expression profiles of all the genes among 10 human tissues, 8 brain regions (amygdala, corpus callosum, cerebellum, caudate nucleus, hippocampus, substania nigra, subthalamic nucleus, and thalamus), spinal cord, fetal brain and fetal liver were also examined by reverse transcription-coupled polymerase chain reaction, products of which were quantified by enzyme-linked immunosorbent assay.

PMID: 10231032, UI: 99246063

Display Abstract Save Text Order Add to Clipboar	rd.
I DISDICE TOURS INTO THE COLOUR FOR	